Between text matter to 2D Code the blank space should be min. 3mm

Position / Orientation shall be changed depending upon the Printer's Machine suitability.

Tacrolimus Capsules, USP

Tacrolimus Capsules, USP
1040242

Space for 15 Digit - Sequential 2D Data matrix Barcode shall be printed by the vendor.

450 mm

750 x 450 mm

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use TACROLIMUS CAPSULES safely and effectively. See full prescribing information for TACROLIMUS CAPSULES.

TACROLIMUS capsules, for oral use.
Initial U.S. Approval: 1994

WARNING: MALIGNANCIES AND SERIOUS INFECTIONS
See full prescribing information for complete boxed warning.

Increased risk for developing serious infections and malignancies with tacrolimus capsules or other immunosuppressants that may lead to hospitalization or death. (5.1, 5.2)

----------INDICATIONS AND USAGE----------

Tacrolimus capsule is a calcineurin-inhibitor immunosuppressant indicated for the prophylaxis of organ rejection in adult and pediatric patients receiving allogeneic liver, kidney or heart transplants, in combination with other immunosuppressants. (1.1)

----------DOSAGE AND ADMINISTRATION----------

Patient Population	Initial Oral Dosage (formulation)	Whole Blood Trough Concentration Range
ADULT		
Kidney Transplant		
With azathioprine or MMF	0.2 mg/kg/day capsules, divided in two doses, every 12 hours	Month 1-3: 7-20 ng/mL, month 4-12: 5-15 ng/mL
With MMF/IL-2 receptor antagonist	0.1 mg/kg/day capsules, divided in two doses, every 12 hours	Month 1-12: 4-11 ng/mL
Liver Transplant		
With corticosteroids only	0.1-0.15 mg/kg/day capsules, divided in two doses, every 12 hours	Month 1-12: 5-20 ng/mL
Heart Transplant		
With azathioprine or MMF	0.075 mg/kg/day capsules, divided in two doses, every 12 hours	Month 1-3: 10-20 ng/mL, Month 4- 6: 5-15 ng/mL
PEDIATRIC		
Kidney Transplant		
	0.3 mg/kg/day capsules or oral suspension, divided in two doses, every 12 hours	Month 1-12: 5-20 ng/mL
Liver Transplant		
	0.15-0.2 mg/kg/ day capsules or 0.2 mg/kg/day oral suspension, divided in two doses, every 12 hours	Month 1-12: 5-20 ng/mL
Heart Transplant		
	0.3 mg/kg/day* capsules or oral suspension, divided in two doses, every 12 hours	Month 1-12: 5-20 ng/mL

MMF = Mycophenolate mofetil
*0.1 mg/kg/day if cell depleting induction treatment is administered.

- Intravenous (IV) use recommended for patients who cannot tolerate oral formulations (capsules or suspension). (2.1, 2.2)
- Administer capsules or suspension consistently with or without food. (2.1)
- Therapeutic drug monitoring is recommended. (2.1, 2.6)
- Avoid eating grapefruit or drinking grapefruit juice. (2.1)
- See dosing adjustments for African-American patients (2.2), hepatic and renal impaired. (2.4, 2.5)

For intravenous dosing information, see the full prescribing information.

----------DOSAGE FORMS AND STRENGTHS----------

- Capsules: 0.5 mg, 1 mg and 5 mg (3)

----------CONTRAINDICATIONS----------

- Hypersensitivity to tacrolimus or HCO-60 (polyoxyl 60 hydrogenated castor oil). (4)

----------WARNINGS AND PRECAUTIONS----------

- Not Interchangeable with Extended Release Tacrolimus Products - Medication Errors: Instruct patients or caregivers to recognize the appearance of tacrolimus capsules. (5.3)
- New Onset Diabetes After Transplant: Monitor blood glucose. (5.4)
- Nephrotoxicity (acute and/or chronic): Reduce the dose, use caution with other nephrotoxic drugs. (5.5)
- Neurotoxicity including risk of Posterior Reversible Encephalopathy Syndrome (PRES): monitor for neurologic abnormalities; reduce or discontinue tacrolimus. (5.6)
- Hyperkalemia: Monitor serum potassium levels. Consider carefully before using with other agents also associated with hyperkalemia. (5.7)
- Hypertension: May require antihypertensive therapy. Monitor relevant drug-drug interactions. (5.8)
- Anaphylactic Reactions with IV formulation: Observe patients receiving tacrolimus injection for signs and symptoms of anaphylaxis. (5.9)
- Not recommended for use with sirolimus: Not recommended in liver and heart transplant due to increased risk of serious infections. (5.10, 7.1)
- Myocardial Hypertrophy: Consider dose reduction/ discontinuation. (5.13)
- Immunizations: Avoid live vaccines. (5.14)
- Pure Red Cell Aplasia: Consider discontinuation of tacrolimus. (5.15)

----------ADVERSE REACTIONS----------

The most common adverse reactions (≥ 15%) were abnormal renal function, hypertension, diabetes mellitus, fever, CMV infection, tremor, hypertrophy, leukopenia, infection, anemia, bronchitis, pericardial effusion, urinary tract infection, constipation, diarrhea, headache, abdominal pain, insomnia, paresthesia, peripheral edema, nausea, hyperkalemia, hypomagnesemia, and hyperglycemia. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Strides Pharma Inc. at 1-877-244-5822 or go to www.strides.com or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----------DRUG INTERACTIONS----------

- Mycophenolate Mofetil Products: Can increase MPA exposure. Interactions from concomitant use with cyclosporine; monitor for MPA-related adverse reactions and adjust MMF or MPA dose as needed. (7.1)
- Sulfasalazine and Grapefruit Juice: Increased tacrolimus concentrations via CYP3A inhibition; avoid concomitant use. (7.2)
- CYP3A Inhibitors: Increased tacrolimus concentrations; monitor concentrations and adjust tacrolimus dose as needed. (5.11, 7.2)
- CYP3A4 Inducers: Decreased tacrolimus concentrations; monitor concentrations and adjust tacrolimus dose as needed. (5.11, 7.2)

----------USE IN SPECIFIC POPULATIONS----------

Pregnancy: Can cause fetal harm. Advise females of reproductive potential to inform healthcare provider of potential risk to the fetus. (8.1, 8.3)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 01/2020

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: MALIGNANCIES AND SERIOUS INFECTIONS

1 INDICATIONS AND USAGE
1.1 Prophylaxis of Organ Rejection in Kidney, Liver, and Heart Transplant

2 DOSAGE AND ADMINISTRATION
2.1 Important Administration Instructions
2.2 Dosing for Adult Kidney, Liver, or Heart Transplant Patients Capsules
2.3 Dosing for Pediatric Kidney, Liver, and Heart Transplant Patients
2.4 Dosage Adjustment in Patients with Renal Impairment
2.5 Dosage Adjustment in Patients with Hepatic Impairment
2.6 Therapeutic Drug Monitoring

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS
5.1 Lymphoma and Other Malignancies
5.2 Serious Infections
5.3 Not Interchangeable With Extended-Release Tacrolimus Products: Medication Errors
5.4 New Onset Diabetes After Transplant
5.5 Nephrotoxicity
5.6 Neurotoxicity
5.7 Hyperkalemia
5.8 Hypertension
5.9 Anaphylactic Reactions with Tacrolimus Injection
5.10 Not Recommended for Use with Sirolimus
5.11 Interactions with CYP3A4 Inhibitors and Inducers
5.12 QT Prolongation
5.13 Myocardial Hypertrophy
5.14 Immunizations
5.15 Pure Red Cell Aplasia

6 ADVERSE REACTIONS ONS
6.1 Clinical Studies Experience
6.2 Postmarketing Adverse Reactions

7 DRUG INTERACTIONS
7.1 Mycophenolic Acid
7.2 Effects of Other Drugs on tacrolimus

FULL PRESCRIBING INFORMATION

WARNING - MALIGNANCIES AND SERIOUS INFECTIONS

Increased risk for developing serious infections and malignancies with tacrolimus or other immunosuppressants that may lead to hospitalization or death. (5.1, 5.2)

1 INDICATIONS AND USAGE

1.1 Prophylaxis of Organ Rejection in Kidney, Liver, and Heart Transplant

Tacrolimus capsules are indicated for the prophylaxis of organ rejection, in adult and pediatric patients receiving allogeneic kidney transplant [*see Clinical Studies (14.1)*], liver transplants [*see Clinical Studies (14.2)*] and heart transplant [*see Clinical Studies (14.3)*], in combination with other immunosuppressants.

2 DOSAGE AND ADMINISTRATION

2.1 Important Administration Instructions

Tacrolimus capsules should not be used without supervision by a physician with experience in immunosuppressive therapy.

Tacrolimus capsules are not interchangeable or substitutable for other tacrolimus extended-release products. This is because rate of absorption following the administration of an extended-release tacrolimus product is not equivalent to that of an immediate-release tacrolimus drug product. Under-or overexposure to tacrolimus may result in graft rejection or other serious adverse reactions. Dosing of immediate-release tacrolimus, modified-release and extended-release dosage forms must occur under physician supervision [*see Warnings and Precautions (5.3)*].

Oral Formulations (Capsules and Oral Suspension)
If patients are able to initiate oral therapy, the recommended starting doses should be initiated. Tacrolimus capsules may be taken with or without food. However, since the presence of food affects the bioavailability of Tacrolimus, if taken with food, it should be taken consistently the same way each time [*see Clinical Pharmacology (12.3)*].

General Administration Instructions
Patients should not eat grapefruit or drink grapefruit juice in combination with Tacrolimus capsules [*see Drug Interactions (7.2)*].

Tacrolimus capsules should not be used simultaneously with cyclosporine. Tacrolimus capsules or cyclosporine should be discontinued at least 24 hours before initiating the other. In the presence of elevated Tacrolimus capsules or cyclosporine concentrations, dosing with the other drug usually should be further delayed.

Therapeutic drug monitoring (TDM) is recommended for all patients receiving Tacrolimus capsules [*see Dosage and Administration (2.6)*].

2.2 Dosing for Adult Kidney, Liver, or Heart Transplant Patients – Capsules

Capsules
If patients are able to tolerate oral therapy, the recommended oral starting doses should be initiated. The initial dose of tacrolimus capsules should be administered no sooner than 6 hours after transplantation in the liver and heart transplant patients. In kidney transplant patients, the initial dose of tacrolimus capsules may be administered within 24 hours of transplantation, but should be delayed until renal function has recovered.

The initial oral tacrolimus capsule dosage recommendations for adult patients with kidney, liver, or heart transplants and whole blood trough concentration range are shown in Table 1. Perform therapeutic drug monitoring (TDM) to ensure that patients are within the ranges listed in Table 1.

Table 1. Summary of Initial Oral Tacrolimus Capsules Dosing Recommendations and Whole Blood Trough Concentration Range in Adults

Patient Population	Tacrolimus Capsules* Initial Oral Dosing	Whole Blood Trough Concentration Range
Kidney Transplant		
With Azathioprine	0.2 mg/kg/day, divided in two doses, administered every 12 hours	Month 1-3: 7-20 ng/mL Month 4-12: 5-15 ng/mL
With MMF/IL-2 receptor antagonist †	0.1 mg/kg/day, divided in two doses, administered every 12 hours	Month 1-12: 4-11 ng/mL
Liver Transplant		
With Corticosteroids Only	0.10-0.15 mg/kg/day, divided in two doses, administered every 12 hours	Month 1-12: 5-20 ng/mL
Heart Transplant		
With azathioprine or MMF	0.075 mg/kg/day, divided in two doses, administered every 12 hours	Month 1-3: 10-20 ng/mL Month 4-12: 5-15 ng/mL

* African-American patients may require higher doses compared to Caucasians [*see Table 2*]

† In a second smaller trial, the initial dose of tacrolimus was 0.15 – 0.2 mg/kg/day and observed tacrolimus concentrations were 6-18 ng/mL during month 1-3 and 5-12 ng/mL during month 4-12 [*see Clinical Studies (14.1)*].

Dosing should be titrated based on clinical assessments of rejection and tolerability. Lower tacrolimus capsule dosages than the recommended initial-dosage may be sufficient as maintenance therapy. Adjunct therapy with adrenal corticosteroids is recommended early post-transplant.

The data in kidney transplant patients indicate that the African-American patients required a higher dose to attain comparable trough concentrations compared to Caucasian patients (Table 2) [*see Use in Specific Populations (8.8) and Clinical Pharmacology (12.3)*].

Table 2. Comparative Dose and Trough Concentration Ratios for Black and Caucasian Patients

Time After Transplant	Caucasian N = 114		African-American N = 56	
	Dose (mg/kg)	Trough Concentrations (ng/mL)	Dose (mg/kg)	Trough Concentrations (ng/mL)
Day 7	0.18	12.2	0.23	10.9
Month 1	0.17	12.8	0.26	12.9
Month 6	0.14	11.8	0.24	11.5
Month 12	0.13	10.1	0.19	11.0

Intravenous Injection
Tacrolimus injection should be used only as a continuous intravenous infusion and should be discontinued as soon as the patient can tolerate oral administration. The first dose of tacrolimus capsules should be given 8-12 hours after discontinuing the intravenous infusion.

The recommended starting dose of tacrolimus injection is 0.03 – 0.05 mg/kg/day in kidney and liver transplant and 0.01 mg/kg/ day in heart transplant given as a continuous intravenous infusion. Adult patients should receive doses at the lower end of the dosing range. Concomitant adrenal corticosteroid therapy is recommended early post-transplantation.

The whole blood trough concentration range described in Table 1 pertains to oral administration of tacrolimus only; while monitoring tacrolimus concentrations in patients receiving tacrolimus injection as a continuous intravenous infusion may have some utility, the observed concentrations will not represent comparable exposures to those evidenced by the trough concentrations observed in patients on oral therapy.

Anaphylactic reactions have occurred with injectables containing castor oil derivatives, such as tacrolimus injection. Therefore, monitoring for signs and symptoms of anaphylaxis is recommended [*see Warnings and Precautions (5.9)*].

2.3 Dosing for Pediatric Kidney, Liver, and Heart Transplant Patients

Oral formulations (capsules)
Pediatric patients in general need higher tacrolimus doses compared to adults: the higher dose requirements may decrease as the child grows older. Recommendations for the initial oral dosing for pediatric transplant patients and whole blood trough concentration range are shown in Table 3. Perform TDM to ensure that patients are within the ranges listed in Table 3.

Table 3. Summary of Initial Tacrolimus Capsule Dosing Recommendations and Whole Blood Trough Concentration Range in Children

Patient Population	Initial Tacrolimus capsule Dosing	Whole Blood Trough Concentration Range
Pediatric kidney transplant patients†	0.3 mg/kg/day capsules or oral suspension, divided in two doses, administered every 12 hours	Month 1-12: 5-20 ng/mL
Pediatric liver transplant patients†	0.15 – 0.2 mg/kg/day capsules or 0.2 mg/kg/day oral suspension, divided in two doses, administered every 12 hours	Month 1-12: 5-20 ng/mL
Pediatric heart transplant patients†	0.3 mg/kg/day* capsules or oral suspension, divided in two doses, administered every 12 hours	Month 1-12: 5-20 ng/mL

* 0.1 mg/kg/day if cell depleting induction treatment is administered.
† *See Clinical Pharmacology (12.2)*

Transition from Oral Tacrolimus to Intravenous Tacrolimus
For conversion of pediatric patients from tacrolimus granules to tacrolimus capsules or from tacrolimus capsules to tacrolimus injection, the same total daily dose should be administered. Monitor tacrolimus blood concentrations and adjust tacrolimus dose as needed. Therapeutic drug monitoring is recommended for all patients receiving Tacrolimus capsules. If a patient is unable to receive an oral formulation, the patient may be started on tacrolimus injection. For pediatric liver transplant patients, the intravenous dose should not exceed 0.03 to 0.05 mg/kg/day.

2.4 Dosage Adjustment in Patients with Renal Impairment

There is potential for nephrotoxicity. Patients should be given to dosing tacrolimus capsules at the lower end of the therapeutic dosing range in patients who have received a liver or heart transplant and have pre-existing renal impairment. Careful monitoring of renal function is necessary. Dose reduction may be required.

In kidney transplant patients with post-operative oliguria, the initial dose of tacrolimus capsules should be administered no sooner than 6 hours and within 24 hours of transplantation, the patient's renal function shows evidence of recovery [*see Dosage and Administration (2.2), Warnings and Precautions (5.5), Use in Specific Populations (8.6), and Clinical Pharmacology (12.3)*].

2.5 Dosage Adjustment in Patients with Hepatic Impairment

Due to the reduced clearance and prolonged half-life, patients with severe hepatic impairment (Child Pugh ≥ 10) may require lower doses of tacrolimus capsules. Close monitoring of blood concentrations is necessary.

Liver transplant patients with severe hepatic impairment experiencing post-transplant hepatic impairment may be associated with increased risk of developing renal insufficiency related to high whole blood concentrations. Careful monitoring of renal function should be monitored closely and dosage adjustments should be considered. Some evidence suggests that lower doses should be used in these patients [*see Dosage and Administration (2.2), Warnings and Precautions (5.5), Use in Specific Populations (8.7) and Clinical Pharmacology (12.3)*].

2.6 Therapeutic Drug Monitoring

Monitoring of tacrolimus blood concentrations in conjunction with other laboratory and clinical parameters is considered an essential aid to patient management for the evaluation of rejection, toxicity, dose adjustments and compliance. Whole blood trough concentration range can be found in Table 1.

Factors influencing frequency of monitoring include but are not limited to hepatic or renal dysfunction, the addition or discontinuation of potentially interacting drugs and the post-transplant time. Blood concentration monitoring is not a replacement for renal and liver function monitoring and biopsies. Data from clinical trials show that tacrolimus whole blood concentrations were most variable during the first week post-transplantation.

The relative risks of toxicity and treatment failure are related to tacrolimus whole blood trough concentrations. Therefore, monitoring of whole blood trough concentrations is recommended to assist in the clinical evaluation of toxicity and efficacy failure.

Methods commonly used for the assay of tacrolimus include high-performance liquid chromatography with tandem mass spectrometric detection (HPLC/MS/MS) and immunoassays. Immunoassays may react with metabolites as well as the parent compound. Whole blood trough concentrations measured using HPLC/MS/MS are lower than those measured by other assays. This bias may depend upon the specific assay and laboratory. Comparison of the concentrations in published literature to patient concentrations using the current assays must be made with detailed knowledge of the assay methods and biological matrices employed. Whole blood is the matrix of choice and specimens should be collected into tubes containing ethylene diamine tetraacetic acid (EDTA) anticoagulant. Heparin anticoagulation is not recommended because of the tendency to form clots on storage. Samples which are not analyzed immediately should be stored at room temperature or in a refrigerator and assayed within 7 days; see assay instructions for specifics. If samples are to be kept longer, they should be deep frozen at -20° C. One study showed drug recovery > 90% for samples stored at -20°C for 6 months, with reduced recovery observed after 6 months.

3 DOSAGE FORMS AND STRENGTHS

Oblong, hard capsule for oral administration contains anhydrous tacrolimus as follows:
- 0.5 mg -opaque red, imprinted "0.5mg" on the cap and "1039" on the body.
- 1 mg -opaque yellow, imprinted "1mg" on the cap and "1039" on the body.
- 5 mg –opaque blue, imprinted "5MG" on the cap and "1039" on the body.

4 CONTRAINDICATIONS

Tacrolimus capsules are contraindicated in patients with a hypersensitivity to tacrolimus. Tacrolimus injection is contraindicated in patients with a hypersensitivity to HCO-60 (polyoxyl 60 hydrogenated castor oil). Anaphylactic/anaphylactoid symptoms reported include dyspnea, rash, pruritus, and acute respiratory distress syndrome [*see Adverse Reactions (6)*].

5 WARNINGS AND PRECAUTIONS

5.1 Lymphoma and Other Malignancies

Patients receiving immunosuppressants, including tacrolimus, are at increased risk of developing lymphomas and other malignancies, particularly of the skin [*see Boxed Warning*]. The risk appears to be related to the intensity and duration of immunosuppression rather than to the use of any specific agent.

As usual for patients with increased risk for skin cancer, examine patients for skin changes; exposure to sunlight and UV light should be limited by wearing protective clothing and using a broad spectrum sunscreen with a high protection factor.

Post-transplant lymphoproliferative disorder (PTLD) has been reported in immunosuppressed organ transplant recipients. The majority of PTLD events appear related to Epstein – Barr Virus (EBV) infection. The risk of PTLD appears greatest in those individuals who are EBV seronegative, a population which includes many young children. Monitor EBV serology during treatment.

5.2 Serious Infections

Patients receiving immunosuppressants, including tacrolimus, are at increased risk of developing bacterial, viral, fungal, and protozoal infections, including opportunistic infections. These infections may lead to serious, including fatal, outcomes. Serious viral infections reported include:
- Polyomavirus associated nephropathy (PVAN), mostly due to BK virus infection
- JC virus-associated progressive multifocal leukoencephalopathy (PML)
- Cytomegalovirus infections: CMV seronegative transplant patients who receive an organ from a CMV seropositive donor disease are at higher risk of developing CMV viremia and CMV disease.

Monitor for the development of infection and adjust the immunosuppressive regimen to balance the risk of rejection with the risk of infection [*see Adverse Reactions (6.1, 6.2)*].

5.3 Not Interchangeable With Extended-Release Tacrolimus Products - Medication Errors

Medication errors, including substitution and dispensing errors, between tacrolimus immediate-release products and tacrolimus extended-release products have been reported. Due to differences in the rate of absorption of tacrolimus, which may result in serious adverse reactions due to under or overexposure to tacrolimus. Tacrolimus is not interchangeable or substitutable for tacrolimus extended-release products. Changes between tacrolimus immediate-release and extended-release dosage forms must occur under physician supervision. Instruct patients and caregivers to recognize the appearance of tacrolimus dosage forms [*see Dosage Forms and Strengths (3)*] and to confirm with the healthcare provider if a different product is dispensed.

5.4 New Onset Diabetes After Transplant

Tacrolimus was shown to cause new onset diabetes mellitus in clinical trials of kidney, liver, and heart transplantation. New onset diabetes after transplantation may be reversible in some patients. African-American and Hispanic kidney transplant patients are at an increased risk. Blood glucose concentrations should be monitored closely in patients using tacrolimus [*see Adverse Reactions (6.1)*].

5.5 Nephrotoxicity

Tacrolimus, like other calcineurin inhibitors, can cause acute or chronic nephrotoxicity. Nephrotoxicity was reported in clinical trials [*see Adverse Reactions (6.1)*]. Consider dosage reduction in patients with elevated serum creatinine or when tacrolimus whole blood trough concentrations greater than the recommended range. The risk for nephrotoxicity may increase when tacrolimus is administered concomitantly with drugs that inhibit CYP3A4 [*see Drug Interactions (7.2)*]. Concomitant use of drugs associated with nephrotoxicity (e.g., aminoglycosides, ganciclovir, amphotericin B, cisplatin, nucleotide reverse transcriptase inhibitors, protease inhibitors) [*see Drug Interactions (7.2)*]. Monitor renal function and consider dosage reduction if nephrotoxicity occurs.

5.6 Neurotoxicity

Tacrolimus may cause a spectrum of neurotoxicities. The most severe neurotoxicities include posterior reversible encephalopathy syndrome (PRES), delirium, seizure and coma; others include tremors, paresthesias, headache, mental status changes, and changes in motor and sensory functions [*see Adverse Reactions (6.1, 6.2)*]. As symptoms may be associated with tacrolimus whole blood trough concentrations at or above the recommended range, monitor for neurologic symptoms and consider dosage reduction or discontinuation of tacrolimus if neurotoxicity occurs.

5.7 Hyperkalemia

Hyperkalemia has been reported with tacrolimus use. Serum potassium levels should be monitored. Careful consideration should be given prior to use of other agents also associated with hyperkalemia (e.g., potassium-sparing diuretics, ACE inhibitors, angiotensin receptor blockers) during tacrolimus therapy [*see Adverse Reactions (6.1)*]. Monitor serum potassium levels periodically during treatment.

5.8 Hypertension

Hypertension is a common adverse effect of tacrolimus therapy and may require antihypertensive therapy [*see Adverse Reactions (6.1)*]. The control of blood pressure can be accomplished with any of the common antihypertensive agents, though careful consideration should be given prior to use of antihypertensive agents associated with hyperkalemia (e.g., potassium-sparing diuretics, ACE inhibitors, angiotensin receptor blockers) [*see Warnings and Precautions (5.7)*]. Calcium-channel blocking agents may increase tacrolimus blood concentrations and therefore require dosage reduction of tacrolimus [*see Drug Interactions (7.2)*].

5.9 Anaphylactic Reactions with Tacrolimus Injection

Anaphylactic reactions have occurred with injectables containing castor oil derivatives, including tacrolimus, in a small percentage of patients (0.6%). The exact cause of these reactions is not known. Tacrolimus injection should be reserved for patients who are unable to take tacrolimus orally. Monitor patients for anaphylaxis when using the intravenous route of administration [*see Dosage and Administration (2.1)*].

5.10 Not Recommended for Use with Sirolimus

Tacrolimus is not recommended for use with sirolimus:
- The use of sirolimus with tacrolimus in studies of de novo liver transplant patients was associated with an excess mortality, graft loss, and hepatic artery thrombosis (HAT) and is not recommended.
- The use of sirolimus (2 mg per day) with tacrolimus in heart transplant patients in a U.S. trial was associated with increased risk of renal function impairment, wound healing complications, and insulin-dependent post-transplant diabetes mellitus, and is not recommended [*see Clinical Studies (14.3)*].

5.11 Interactions with CYP3A4 Inhibitors and Inducers

When co-administering tacrolimus with strong CYP3A4 inhibitors (e.g., diltiazem, bromocriptine, clotrimazole, ketoconazole, itraconazole, nicardipine, nifedipine, verapamil, fluconazole, voriconazole, erythromycin, clarithromycin, telithromycin, and nefazodone), antibiotics in the dosing regimen of tacrolimus and subsequent frequent monitoring of tacrolimus whole blood trough concentrations and tacrolimus-associated adverse reactions are recommended [*see Drug Interactions (7)*].

5.12 QT Prolongation

Tacrolimus may prolong the QT/QTc interval and may cause Torsade de Pointes. Avoid tacrolimus in patients with congenital long QT syndrome. In patients with congestive heart failure, bradyarrhythmias, those taking certain antiarrhythmic medications or other medical products that lead to QT prolongation, and those with electrolyte disturbances such as hypokalemia, hypocalcemia, or hypomagnesemia, consider obtaining electrocardiograms and monitoring electrolytes periodically. If possible, correct any electrolyte during treatment.

When co-administering tacrolimus with other substrates and/or inhibitors of CYP3A4 that also have the potential to prolong the QT interval, a reduction in tacrolimus dose, frequent monitoring of tacrolimus whole blood concentrations, and monitoring for QT prolongation is recommended [*see Drug Interactions (7.2)*]. Tacrolimus can also cause hypomagnesemia, which can result in increased tacrolimus whole blood concentrations with or without concurrent QT prolongation [*see Drug Interactions (7.2)*].

5.13 Myocardial Hypertrophy

Myocardial hypertrophy has been reported in infants, children, and adults, particularly those with high tacrolimus trough concentrations, and is generally manifested by echocardiographically demonstrated concentric increases in left ventricular posterior wall and interventricular septum thickness. This condition appears reversible in most cases following dose reduction or discontinuance of therapy. In patients who develop renal failure or clinical manifestations of ventricular dysfunction while receiving tacrolimus therapy, echocardiographic evaluation should be considered. If myocardial hypertrophy is diagnosed, dosage reduction or discontinuation of tacrolimus should be considered [*see Adverse Reactions (6.2)*].

5.14 Immunizations

Whenever possible, administer the complete complement of vaccines before transplantation and treatment with tacrolimus.

The use of live vaccines should be avoided during treatment with tacrolimus; examples include (not limited to) the following: intranasal influenza, measles, mumps, rubella, oral polio, BCG, yellow fever, varicella, and TY21a typhoid vaccines.

Inactivated vaccines noted to be safe for administration after transplantation may not be sufficiently immunogenic during treatment with tacrolimus.

5.15 Pure Red Cell Aplasia

Cases of pure red cell aplasia (PRCA) have been reported in patients treated with tacrolimus. A mechanism for tacrolimus-induced PRCA has not been elucidated. All patients reported risk factors for PRCA such as parvovirus B19 infection, underlying disease, or concomitant medications associated with PRCA. If PRCA is diagnosed, discontinuation of tacrolimus should be considered [*see Adverse Reactions (6.2)*].

6 ADVERSE REACTIONS

6.1 Clinical Studies Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice. In addition, the clinical trials were not designed to establish comparative differences across study arms with regards to the adverse reactions discussed below.

Kidney Transplantation
The incidence of adverse reactions was determined in three randomized kidney transplant trials. One of the trials used azathioprine (AZA) and corticosteroids and two of the trials used mycophenolate mofetil (MMF) and corticosteroids concomitantly for maintenance immunosuppression.

The most common adverse reactions in conjunction with azathioprine and corticosteroids following kidney transplantation was assessed in a trial were 205 patients received Tacrolimus-based immunosuppression and 207 patients following received cyclosporine-based immunosuppression. The trial population had a mean age of 43 years (mean ± SD was 43 ± 13 years on Tacrolimus and 44 ± 12 years on cyclosporine arm), the distribution was 61% male, and the composition was White (58%), African-American (25%), Asian (7%), Hispanic (11%), and Other (3%). The 12-month post-transplant adverse reactions are presented below.

The most common adverse reactions (≥ 30%) observed in tacrolimus-treated kidney transplant patients are: infection, tremor, hypertrophy, diarrhea, hypertension, nausea, abnormal renal function, abdominal pain, insomnia, paresthesia, anemia, pain, fever, asthenia, constipation, hypoglycemia, peripheral edema, and dyspepsia. Adverse reactions reported in approximately 5% of kidney transplant patients receiving tacrolimus in the U.S. and European randomized trials.

Table 4. Kidney Transplantation: Adverse Reactions Occurring in ≥ 15% of Patients Treated with Tacrolimus in Conjunction with Azathioprine (AZA)

	Tacrolimus /AZA (N = 205)	Cyclosporine/AZA (N = 207)
Nervous System		
Tremor	54%	34%
Headache	44%	38%
Insomnia	32%	30%
Paresthesia	23%	16%
Dizziness	19%	16%
Gastrointestinal		
Diarrhea	44%	41%
Nausea	38%	36%
Constipation	35%	43%
Vomiting	29%	23%
Dyspepsia	28%	20%
Cardiovascular		
Hypertension	50%	52%
Chest Pain	19%	13%
Urogenital		
Creatinine Increased	45%	42%
Urinary Tract Infection	34%	35%
Metabolic and Nutritional		
Hypomagnesemia	49%	53%
Hyperglycaemia	34%	17%
Hyperlipidemia	31%	38%
Diabetes Mellitus	24%	9%
Hyperkalemia	22%	25%
Hypophosphatemia	22%	16%
Edema	18%	19%
Hemic and Lymphatic		
Anemia	30%	24%
Leukopenia	15%	17%
Miscellaneous		
Infection	45%	49%
Peripheral Edema	38%	48%
Asthenia	34%	30%
Abdominal Pain	33%	31%
Pain	32%	30%
Fever	29%	29%
Back Pain	24%	20%
Respiratory System		
Dyspnea	22%	18%
Cough Increased	18%	15%
Musculoskeletal		
Arthralgia	25%	24%
Skin		
Rash	17%	12%
Pruritus	15%	7%

Two trials were conducted for tacrolimus-based immunosuppression in conjunction with MMF and corticosteroids. In the non-US trial (Study 1), the incidence of adverse reactions was based on 1195 kidney transplant patients that received tacrolimus (Group A) or cyclosporine (Group B) in conjunction with MMF in Study 1. All patients received a regimen of MMF and corticosteroids; all patients, except those in one of the two cyclosporine groups, also received induction with daclizumab. The trial population had a mean age of 46 years (range 17 to 76); the distribution was 65% male, and the composition was 93% Caucasian. The 12 month post-transplant information from this trial is presented below.

Adverse reactions that occurred in ≥ 10% of kidney transplant patients treated with tacrolimus in conjunction with MMF in Study 1 (Note: This trial was conducted entirely outside of the United States. Such trials often report a lower incidence of adverse reactions in comparison to U.S. trials) are presented below:

Table 5. Kidney Transplantation: Adverse Reactions Occurring in ≥10% of Patients Treated with Tacrolimus in Conjunction with MMF (Study 1)

	Tacrolimus (Group C) (N=403)	Cyclosporine (Group A) (N=384)	Cyclosporine (Group B) (N=408)
Diarrhea	25%	16%	13%
Urinary Tract Infection	24%	28%	24%
Anemia	17%	19%	17%
Hypertension	13%	14%	12%
Leukopenia	13%	10%	10%
Edema peripheral	11%	12%	13%
Hyperlipidemia	10%	15%	13%

Key: Group A = OsA/MMF/CS, B = OsA/MMF/CS/Daclizumab, C = Tac/MMF/CS/Daclizumab
OsA = Cyclosporine, CS = Corticosteroids, Tac = Tacrolimus, MMF = mycophenolate mofetil

In the U.S. trial (Study 2) with tacrolimus-based immunosuppression in conjunction with MMF and corticosteroids, 424 kidney transplant patients received tacrolimus (n=212) or cyclosporine (n=212) in combination with MMF 1 gram twice daily, basiliximab induction, and corticosteroids. The trial population had a mean age of 49 years (range 17 to 77); the distribution was 63% male, and the composition was White (74%), African-American (20%), Asian (3%), and Other (3%). The 12-month post-transplant information from this trial is presented below.

Adverse reactions that occurred in ≥ 15% of kidney transplant patients treated with tacrolimus in conjunction with MMF in Study 2 are presented below:

Table 6. Kidney Transplantation: Adverse Reactions Occurring in ≥ 15% of Patients Treated with Tacrolimus in Conjunction with MMF (Study 2)

	Tacrolimus/MMF (N =212)	Cyclosporine/MMF (N =212)
Gastrointestinal Disorders		
Diarrhea	44%	26%
Nausea	39%	47%
Constipation	36%	41%
Vomiting	26%	25%
Dyspepsia	18%	15%
Injury, Poisoning, and Procedural Complications		
Post-Procedural Pain	29%	27%
Graft Dysfunction	28%	23%
Graft Dysfunction	24%	18%
Metabolism and Nutrition Disorders		
Hypomagnesemia	28%	22%
Hypophosphatemia	26%	21%
Hyperkalemia	26%	19%
Hyperglycemia	21%	15%
Hypercholesterolemia	18%	25%
Hypokalemia	16%	18%
Nervous System Disorders		
Tremor	34%	20%
Headache	24%	25%
Blood and Lymphatic System Disorders		
Anemia	30%	28%
Leukopenia	16%	12%
Miscellaneous		
Edema Peripheral	35%	46%
Hypertension	32%	35%
Insomnia	30%	21%
Urinary Tract Infection	26%	22%
Blood Creatinine Increased	23%	23%

Less frequently observed adverse reactions in kidney transplantation patients are described under the subsection "Less Frequently Reported Adverse Reactions (> 2% and < 15%) in Liver, Kidney, and Heart Transplant Studies."

Liver Transplantation
The two large randomized comparative liver transplant trials. In the U.S. trial, 263 adult and pediatric patients received tacrolimus and steroids and 266 patients received cyclosporine-based immunosuppression regimen (CsA/AZA). The trial population had a mean age of 52 years (range 18 to 70). The distribution was 62% male, and the composition was White (78%), African-American (5%), Asian (2%), Hispanic (13%), and Other (2%). In the European trial, 270 patients received tacrolimus and steroids and 275 patients received CsA/AZA. The trial population had a mean age of 46 years (range 13 to 68); the distribution was 59% male, and the composition was White (95.4%), Black (1%), Asian (2%), and Other (2%).

The proportion of patients reporting more than one adverse event was > 99% in both the tacrolimus group and the CsA/AZA group. Precautions must be taken when comparing the incidence of adverse reactions in the U.S. trial to that in the European trial. The 12-month post-transplant information from the U.S. trial and from the European trial is presented below. The two trials also included different patient populations and patients were treated with immunosuppressive regimens of differing intensities. Adverse reactions reported in ≥ 15% in tacrolimus patients (combined trial results) are presented below for the two controlled

type of tacrolimus product. **Tacrolimus extended-release medicines are not the same as tacrolimus capsules or granules** and cannot be substituted for each other, unless specifically prescribed by your healthcare provider, who will send you to get blood tacrolimus levels at a lab. **Check your tacrolimus when you get a new prescription and before you take it to make sure you have received tacrolimus capsule or tacrolimus granules.**

- Check with the pharmacist and call your healthcare provider if you think you were given the wrong medicine.
- **high blood sugar (diabetes).** Your healthcare provider may do blood tests to check for diabetes while you take tacrolimus capsules. Call your healthcare provider right away if you have any symptoms of high blood sugar, including:
  - o frequent urination
  - o increased thirst or hunger
  - o blurred vision
  - o confusion
  - o drowsiness
  - o loss of appetite
  - o fruity smell on your breath
  - o nausea, vomiting, or stomach pain
- **kidney problems.** Kidney problems are a serious and common side effect of tacrolimus. Your healthcare provider may do blood tests to check your kidney function while you take tacrolimus capsules.
- **nervous system problems.** Nervous system problems are a serious and common side effect of tacrolimus. Call your healthcare provider right away if you get any of these symptoms while taking tacrolimus capsules. These could be signs of a serious nervous system problem:
  - o headache
  - o confusion
  - o seizures
  - o changes in your vision
  - o changes in behavior
  - o coma
  - o tremors
  - o numbness and tingling
- **high levels of potassium in your blood.** Your healthcare provider may do blood tests to check your potassium level while you take tacrolimus capsules.
- **high blood pressure.** High blood pressure is a serious and common side effect of tacrolimus. Your healthcare provider will monitor your blood pressure while you take tacrolimus and may prescribe blood pressure medicine for you, if needed. Your healthcare provider may instruct you to check your blood pressure at home.
- **changes in the electrical activity of your heart (QT prolongation).**
- **heart problems (myocardial hypertrophy).** Tell your healthcare provider right away if you get any of these symptoms of heart problems while taking tacrolimus capsules:
  - o shortness of breath
  - o chest pain
  - o feel lightheaded
  - o feel faint
- **severe low red blood cell count (anemia).**

The most common side effects of tacrolimus capsules in people who have received a kidney, liver or heart transplant are:

- infections in general, including cytomegalovirus (CMV) infection
- tremors (shaking of the body)
- constipation
- diarrhea
- headache
- stomach pain
- trouble sleeping
- nausea
- high blood sugar (diabetes)
- low levels of magnesium in your blood
- low levels of phosphate in your blood
- swelling of the hands, legs, ankles, or feet
- weakness
- pain
- high levels of fat in your blood
- high levels of potassium in your blood
- low red blood cell count (anemia)
- low white blood cell count
- fever
- numbness or tingling in your hands and feet
- inflammation of your airway (bronchitis)
- fluid around your heart

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of tacrolimus capsules. For more information, ask your healthcare provider or pharmacist.

Call your healthcare provider for medical advice about side effects. You may report side effects to **Strides Pharma Inc at 1-877-244-9825 or go to www.strides.com or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch**

How should I store tacrolimus capsules?
Store Tacrolimus capsules at room temperature between 68°F to 77°F (20°C-25°C)

Keep tacrolimus capsules and all medicines out of the reach of children.

General information about the safe and effective use of tacrolimus capsules

- Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use tacrolimus capsules for a condition for which it was not prescribed. Do not give tacrolimus capsules to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about tacrolimus that is written for health professionals
- This Patient Information leaflet summarizes the most important information about Tacrolimus. If you would like more information, talk to your healthcare provider.

What are the ingredients in tacrolimus capsules?
Active ingredient: tacrolimus, USP

Inactive ingredients: croscarmellose sodium NF, hypromellose USP, lactose monohydrate NF, and magnesium stearate NF. The 0.5 mg capsule shell contains, FD & C Blue 1, FD & C Red 40, gelatin NF and titanium dioxide USP; the 1 mg capsule shell contains D & C Red 28, D & C Yellow 10, FD & C Blue 1, gelatin NF, and titanium dioxide USP.

Manufactured by:
Strides Pharma Science Ltd.
Bengaluru - 562106, India

Distributed by:
Strides Pharma Inc.
East Brunswick, NJ 08816

Revised: 01/2020

This Patient Information has been approved by the U.S. Food and Drug Administration.

1040242

459 mm

- Gastrointestinal: Bile duct stenosis, colitis, enterocolitis, gastrointestinitis, gastrosophageal reflux disease, hepatic cytolysis, hepatic necrosis, hepatobiliary, impaired gastric emptying, liver fatty, mouth ulceration, pancreatitis hemorrhagic, pancreatitis necrotizing, stomach ulcer, veno-occlusive liver disease
- Hemic/Lymphatic: Agranulocytosis, disseminated intravascular coagulation, hemolytic anemia, neutropenia, febrile neutropenia, pancytopenia, thrombocytopenic purpura, thrombotic thrombocytopenic purpura, pure red cell aplasia *[see Warnings and Precautions (5.15)]*
- Infections: Cases of progressive multifocal leukoencephalopathy (PML), sometimes fatal; -polyoma virus-associated-nephropathy (PVAN) including graft loss *[see Warnings and Precautions (5.2)]*
- Metabolic/Nutritional: Glycosuria, increased amylase including pancreatitis, weight decreased
- Miscellaneous: Feeling hot and cold, feeling jittery, hot flushes, multi-organ failure, primary graft dysfunction
- Musculoskeletal and Connective Tissue Disorders: Pain in extremity including Calcineurin-Inhibitor Induced Pain Syndrome (CIPS)
- Nervous System: Carpal tunnel syndrome, cerebral infarction, hemiparesis, leukoencephalopathy, mental disorder, mutism, nervous system hemorrhages including cerebral hemorrhage, CNS vascular disorder, PRES (Posterior reversible encephalopathy syndrome), *[Warnings and Precautions (5.2)]*, quadriplegia, speech disorder, syncope
- Respiratory: Acute respiratory distress syndrome, interstitial lung disease, lung infiltration, respiratory distress, respiratory failure
- Skin: Stevens-Johnson syndrome; toxic epidermal necrolysis
- Special Senses: Blindness, optic neuropathy, tinnitus-cortical, hearing loss including deafness, photophobia
- Urogenital: Acute renal failure, cystitis hemorrhagic, hemolytic-uremic syndrome

7 DRUG INTERACTIONS

7.1 Concomitant Medications

When tacrolimus is prescribed with a given dose of a mycophenolic acid (MPA) product, exposure to MPA is higher with tacrolimus co-administration than with cyclosporine co-administration with MPA, because cyclosporine interrupts the enterohepatic recirculation of MPA while tacrolimus does not. Monitor for MPA-associated adverse reactions and reduce the dose of concomitantly administered mycophenolic acid products as needed.

7.2 Effects of Other Drugs on Tacrolimus

Table 15 displays the effects of other drugs on Tacrolimus

Table 15: Effects of Other Drugs on Tacrolimus Substances in Tacrolimus*

Drug/Substance Class or Name	Drug Interaction Effect	Recommendations
Grapefruit or grapefruit juice†	May increase tacrolimus whole blood trough concentrations and increase the risk of serious adverse reactions (e.g., neurotoxicity, QT prolongation) *[see Warnings and Precautions (5.6, 5.7,11.5,12)]*.	Avoid grapefruit or grapefruit juice.
Strong CYP3A Inducers‡: Antimycobacterials (e.g., rifampin, rifabutin), anticonvulsants (e.g., phenytoin, carbamazepine and phenobarbital), St. John's Wort	Increase tacrolimus dose and monitor tacrolimus whole blood trough concentrations and increase the risk of rejection *[see Warnings and Precautions (5.11)]*.	Increase tacrolimus dose and monitor tacrolimus whole blood trough concentrations and increase the risk of rejection *[see Drug Interactions (7.2, 2.6) and Clinical Pharmacology (12.3)]*.
Strong CYP3A Inhibitors‡: Protease inhibitors (e.g., nelfinavir, telaprevir, boceprevir, ritonavir), azole antifungals (e.g., voriconazole, posaconazole, itraconazole, clotrimazole, ketoconazole), erythromycin, telithromycin, clarithromycin, nefazodone, telaprevir, cobicistat-containing regimens	May increase tacrolimus whole blood trough concentrations and increase the risk of serious adverse reactions (e.g., neurotoxicity, QT prolongation) *[see Warnings and Precautions (5.6, 5.7,11.5,12)]*.	Reduce tacrolimus dose (for voriconazole and posaconazole, see prescribing information) and adjust dose based on tacrolimus whole blood trough concentrations *[see Dosage and Administration (2.6) and Clinical Pharmacology (12.3)]*.
Mild or Moderate CYP3A Inducers: Glucocorticoids, antibiotics (e.g., erythromycin, sulfamethoxazole and trimethoprim), calcium channel blockers (e.g., verapamil, diltiazem, nifedipine), nafcillin, isoniazid, amiodarone, danazol, ethinyl estradiol, chloramphenicol, metamizole and omeprazole	May increase tacrolimus whole blood trough concentrations and increase the risk of serious adverse reactions (e.g., neurotoxicity, QT prolongation) *[see Warnings and Precautions (5.6, 5.7,11.5,12)]*.	Monitor tacrolimus whole blood trough concentrations and reduce tacrolimus dose if needed *[see Dosage and Administration (2.6) and Clinical Pharmacology (12.3)]*.
Other drugs, such as: Magnesium and aluminum hydroxide antacids, Metoclopramide	May increase tacrolimus whole blood trough concentrations and increase the risk of serious adverse reactions (e.g., neurotoxicity, QT prolongation) *[see Warnings and Precautions (5.6, 5.7,11.5,12)]*.	Monitor tacrolimus whole blood trough concentrations and reduce tacrolimus dose if needed *[see Dosage and Administration (2.6) and Clinical Pharmacology (12.3)]*.
Mild or Moderate CYP3A Inducers: Methylprednisolone, prednisone	May decrease tacrolimus concentrations.	Monitor tacrolimus whole blood trough concentrations and adjust tacrolimus dose if needed *[see Dosage and Administration (2.6)]*.

*Tacrolimus dosage adjustment recommendation based on observed effect of coadministered drug on tacrolimus exposure *[see Clinical Pharmacology (12.3)]*, literature reports of altered tacrolimus exposures, or the other drug's known CYP3A inhibitory/ inducer status.

†High dose or double strength grapefruit juice is a strong CYP3A inhibitor; low dose or single strength grapefruit juice is a moderate CYP3A inhibitor.

‡Strong CYP3A inhibitor/inducer, based on reported effect on exposures to tacrolimus along with supporting *in vitro* CYP3A inhibition/induction data, or based on drug-drug interaction studies with midazolam (sensitive CYP3A probe substrate).

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Exposure Registry

There is an ongoing registry that monitors pregnancy outcomes in women exposed to tacrolimus during pregnancy. The Transplantation Pregnancy Registry International (TPRI) is a voluntary pregnancy exposure registry that monitors outcomes of pregnancy in female transplant recipients and their fetuses fathered by male transplant recipients exposed to immunosuppressants including tacrolimus. Healthcare providers are encouraged to advise their patients to register by contacting the Transplantation Pregnancy Registry International at 877-955-6837 or https://www.transplantpregnancyregistry.org/.

Risk Summary

Tacrolimus can cause fetal harm when administered to a pregnant woman. Data from postmarketing surveillance and TPRI suggest that infants exposed to tacrolimus *in utero* are at a risk of prematurity, birth defects/congenital anomalies, low birth weight, and fetal distress *[see Human Data]*. Advise pregnant women of the potential risk to the fetus.

Administration of oral tacrolimus to pregnant rabbits and rats throughout the period of organogenesis was associated with maternal toxicity/lethality, as well as increased incidence of fetal malformations, malformations and embryo/fetal death at clinically relevant doses (0.3 to 6.9 times the recommended clinical dose (0.2 to 0.075 mg/kg/day), on a mg/m² basis). Administration of oral tacrolimus to rats prior to mating, and throughout gestation and lactation produced maternal toxicity/lethality, embryo/fetal toxicity and reduced pup viability at clinically relevant doses (0.8 to 6.9 times the recommended clinical dose range, on a mg/m² basis). Intravenous/ipsilateral septal defects, hydronephrosis, craniofacial malformations and skeletal effects were observed in offspring that died. Effects on parturition (incomplete delivery of nonviable pups) were observed at 1 mg/kg (0.8 to 2.2 times the recommended clinical dose range) *[see Nonclinical Toxicology (13.1)]*.

8.2 Lactation

Risk Summary

Controlled lactation studies have not been conducted in humans; however, tacrolimus has been reported to be present in human milk. The effects of tacrolimus on the breastfed infant, or on milk production have not been assessed. Tacrolimus is excreted in rat milk and in peri-/postnatal rat studies; exposure to tacrolimus during the postnatal period was associated with developmental toxicity in the offspring at clinically relevant doses *[see Pregnancy (8.1) and Nonclinical Toxicology (13.1)]*.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for tacrolimus and any potential adverse effects on the breastfed child from tacrolimus or from the underlying maternal condition.

8.3 Females and Males of Reproductive Potential

Contraception

Tacrolimus can cause fetal harm when administered to pregnant women. Advise female and male patients of reproductive potential to speak to their healthcare provider on family planning options before using appropriate contraception prior to taking tacrolimus, during tacrolimus therapy, and for 4 weeks after stopping tacrolimus therapy.

Infertility

Based on findings in animals, male and female fertility may be compromised by treatment with tacrolimus *[see Nonclinical Toxicology (13.1)]*.

8.4 Pediatric Use

Safety and effectiveness have been established in pediatric liver, kidney, and heart transplant patients.

Liver transplant

Safety and efficacy using tacrolimus granules in pediatric de novo liver transplant patients less than 16 years of age are based on a randomized, active controlled study that included 56 pediatric patients, 37 of which received tacrolimus granules, and supported by two pharmacokinetic and safety studies in 151 children who received tacrolimus. Additionally, 122 pediatric patients were studied in an uncontrolled trial of tacrolimus in living related donor liver transplantation. Dose adjustments were made in the PK studies based on clinical status and whole blood concentrations. Pediatric patients generally required higher doses of tacrolimus to maintain blood concentrations comparable to adult patients *[see Dosage and Administration (2.2), Adverse Reactions (6.1), Clinical Pharmacology (12.3) and Clinical Studies (14.2)]*.

Kidney transplant

Safety and efficacy in pediatric and tacrolimus granules in pediatric kidney and heart transplant patients is supported by adequate and well-controlled studies and pharmacokinetic data in adult kidney and heart transplant patients with additional pharmacokinetic data in pediatric liver and heart transplant patients and safety data in pediatric liver transplant patients *[see Dosage and Administration (2.3) and Clinical Pharmacology (12.3)]*.

8.5 Geriatric Use

Clinical trials of tacrolimus did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

8.6 Renal Impairment

The pharmacokinetics of tacrolimus in patients with renal impairment was similar to that in healthy volunteers with normal renal function. However, consideration should be given to dosing tacrolimus at the lower end of the therapeutic dosing range in patients with severe renal impairment. In heart transplant and liver post-transplant renal impairment, further reductions in dose below the dosing range may be required *[see Dosage and Administration (2.4) and Clinical Pharmacology (12.3)]*.

8.7 Hepatic Impairment

The mean clearance of tacrolimus was substantially lower in patients with severe hepatic impairment (mean Child-Pugh score: >10) compared to healthy volunteers with normal hepatic function. Close monitoring of tacrolimus trough concentrations is warranted in patients with hepatic impairment *[see Clinical Pharmacology (12.3)]*.

The use of tacrolimus in liver transplant recipients experiencing post-transplant hepatic impairment may be associated with increased risk of developing renal insufficiency relative to liver transplant recipients without hepatic impairment. These patients should be monitored closely and dosage adjustments should be considered. Some evidence suggests that lower doses should be used in these patients *[see Dosage and Administration (2.5) and Clinical Pharmacology (12.3)]*.

8.8 Race or Ethnicity

African-American patients may need to be titrated to higher dosages to attain comparable trough concentrations compared to Caucasian patients *[see Dosage and Administration (2.2) and Clinical Pharmacology (12.3)]*.

Limited data from transplant patients at increased risk for rare small bowel diarrhea after transplant. Monitor blood glucose concentrations and treat appropriately *[see Warnings and Precautions (5.4)]*.

10 OVERDOSAGE

Limited overdosage experience is available. Acute overdosages of up to 30 times the intended dose have been reported. Almost all cases have resolved without sequelae and all patients recovered with no sequelae. Acute overdosage was sometimes followed by adverse reactions consistent with those listed in *Adverse Reactions (6)* (including tremors, abnormal renal function, hypertension, infections, convulsions, and QT prolongation). Gastrointestinal symptoms (including nausea, vomiting, and diarrhea), electrolyte imbalance, and serious cardiac and/or neurological disorders (including seizures, arrhythmias, and coma) have also been reported. Should overdosage occur, it is anticipated that tacrolimus is not dialyzable to any significant extent; there is no experience with charcoal hemoperfusion. The oral use of activated charcoal has been reported in treating acute overdose, but experience to date has been insufficient to warrant recommending its use. General supportive measures and treatment of specific symptoms should be followed in all cases of overdosage.

In acute oral and IV toxicity studies, mortalities were seen at or above the following doses: in adult rats, 52 times the recommended human oral dose; in immature rats, 16 times the recommended oral dose; and in adult rats, 16 times the recommended human IV dose (all based on body surface area corrections).

11 DESCRIPTION

Tacrolimus, previously known as FK506, is the active ingredient in Tacrolimus capsules. Tacrolimus is a calcineurin-inhibitor immunosuppressant. Tacrolimus has the chemical name [3*S*-[3*R**[*E*(1*S**,3*S**,4*S**)],4*S**,5*R**,8*S**,9*E*,12*R**,14*R**,15*S**,16*R**,18*S**,19*S**,26*aR**]]-5,6,8,11,12,13,14,15,16, 17,18,19,24,25,26,26*a*-hexadecahydro-5,19-dihydroxy-3-[2-(4-hydroxy-3-methoxycyclohexyl)-1-methylethenyl]-14, 16-dimethoxy-4,10,12,18-tetramethyl-8-(2-propenyl)-15,19-epoxy-3*H*-pyrido[2,1-*c*][1,4]oxaazacyclotricosine-1,7,20,21 (4*H*,23*H*)-tetrone, monohydrate.

The chemical structure of tacrolimus is:

Tacrolimus has an empirical formula of $C_{44}H_{69}NO_{12}$•H_2O and a formula weight of 822.03. Tacrolimus appears as white crystals or crystalline powder. It is practically insoluble in water, freely soluble in ethanol, and very soluble in methanol and chloroform.

Tacrolimus is available for oral administration as capsules (tacrolimus capsules USP) containing either the equivalent of 0.5 mg, 1 mg or 5 mg of anhydrous tacrolimus USP. Inactive ingredients include croscarmellose sodium NF, hypromellose USP, lactose monohydrate NF, and magnesium stearate NF. The 0.5 mg capsule shell contains D & C Blue 1, FD & C Red 40, gelatin NF and titanium dioxide USP; the 1 mg capsule shell contains FD & C Blue 1, gelatin NF, iron oxide yellow and titanium dioxide USP; and the 5 mg capsule shell contains D & C Red 28, D & C Yellow 10, FD & C Blue 1, gelatin NF, and titanium dioxide USP.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Tacrolimus binds to an intracellular protein, FKBP-12. A complex of tacrolimus-FKBP-12, calcium, calmodulin, and calcineurin is then formed. The immunosuppressant and other actions of tacrolimus may result from the phosphatase activity of calcineurin is inhibited. Such inhibition prevents the dephosphorylation and translocation of various factors such as the nuclear factor of activated T-cells (NF-AT), and nuclear factor kappa-light-chain-enhancer of activated B-cells (NF-κB).

Tacrolimus inhibits the expression and/or production of several cytokines that include interleukin (IL)-1 beta, IL-2, IL-3, IL-4, IL-5, IL-6, IL-8, IL-10, tumor necrosis factor (TNF)-alpha, granulocyte-macrophage colony-stimulating factor (GM-CSF), and interferon (IFN)-gamma. Tacrolimus also inhibits IL-2 receptor expression and nitric oxide release, and the proliferation of transforming growth factor beta that can lead to immunosuppressive activity. The net result is the inhibition of T-lymphocyte activation and proliferation, but the exact mechanism of tacrolimus' effect on transplantation is unknown.

12.2 Pharmacokinetics

The pharmacokinetic profile of tacrolimus is dependent on the patient's clinical status. Tacrolimus is extensively bound to plasma proteins, binding primarily to albumin and alpha-1-acid glycoprotein, and has a high level of association with erythrocytes. The distribution of tacrolimus between plasma and erythrocytes is dependent on several factors, such as hematocrit, temperature and the albumin and alpha-1-acid glycoprotein concentrations. Whole blood and plasma concentrations are approximately 77 and 33 ng/mL, respectively, after administration of plasma separation, drug concentration, and plasma protein concentration. In a U.S. trial, the ratio of whole blood concentration to plasma concentration averaged 20 (range 13 to 91).

Elimination

Tacrolimus is extensively metabolized by the mixed-function oxidase system, primarily the cytochrome P-450 system (CYP3A). A metabolic pathway leading to the formation of 8 possible metabolites has been proposed. Demethylation and hydroxylation were identified as the primary mechanisms of biotransformation *in vitro*. The major metabolite identified in incubations with human liver microsomes is 13-demethyl tacrolimus. *In vitro* studies, a 31-demethyl metabolite has been reported to have the same activity as tacrolimus.

Excretion

The mean clearance following IV administration of tacrolimus is 0.040, 0.083, 0.053 and 0.051 L/hr/kg in healthy volunteers, adult kidney transplant patients, adult liver transplant patients, and adult heart transplant patients, respectively; in man, less than 1% of the administered dose is excreted unchanged in urine.

In a mass balance study of 8 radiolabeled radiolabeled tacrolimus to 6 healthy volunteers, the mean recovery of radiolabel was 93.6% ± 1.36% (SD) over 96 hours, with 2.07% ± 0.39% recovered in urine and 91.5% ± 1.39% recovered in feces. The 13.9 hours whereas it was 43.3 ± 11.4 hours based on tacrolimus concentrations. The mean dose recovered in bile was 4.28% ± 0.015 L/hr/kg and clearance of tacrolimus was 0.029 ± 0.006 L/hr/kg. When administered IPO, the mean recovery of the elimination half-life based on radioactivity was 31.9 ± 10.5 hours whereas it was 48.4 ± 12.3 hours based on tacrolimus concentrations. The mean clearance of radiolabel was 0.226 ± 0.116 L/hr/kg and clearance of tacrolimus 0.172 ± 0.066 L/hr/kg.

Specific Populations

Pediatric Patients

Tacrolimus capsules Pharmacokinetics in Pediatric Patients

Pharmacokinetics of tacrolimus have also been studied in liver transplantation patients, 0.7 to 13.2 years of age. Following IV administration of 0.037 mg/kg/day (0.5 to 2 pediatric patients (8 male and 4 female), mean terminal half-life and clearance were 19.2 ± 5.0 hours and 0.12 ± 0.04 L/hr/kg, respectively. Following oral administration to the same patients, mean AUC and C_{max} were 181 ± 65 ng•hr/mL and 30 ± 11 ng/mL, respectively. The absolute bioavailability was 19 ± 24%.

Pharmacokinetics of tacrolimus have also been studied in kidney transplantation patients, 8.2 ± 2.4 years of age. Following IV infusion of a 0.06 mg/kg/day to 12 pediatric patients (8 male and 4 female), mean terminal half-life and clearance were 19.2 ± 5.0 hours and 0.12 ± 0.04 L/hr/kg, respectively. Following oral administration to the same patients, mean AUC and C_{max} were 181 ± 65 ng•hr/mL and 30 ± 11 ng/mL, respectively. The absolute bioavailability was 19 ± 24%.

Whole blood trough concentrations from 31 patients less than 12 years old showed that pediatric patients needed higher doses than adults to achieve similar tacrolimus trough concentrations *[see Dosage and Administration (2.2)]*.

Renal and Hepatic Impaired Patients

The mean pharmacokinetic parameters for tacrolimus following single administrations to adult patients with renal and hepatic impairment are given in Table 19.

Table 19. Pharmacokinetics in Renal and Hepatic Impaired Adult Patients

Population (No. of Patients)	Dose	$AUC_{0-\infty}$ (ng•hr/mL)	$t_{1/2}$ (hr)	V (L/kg)	Cl (L/hr/kg)
Renal Impairment (n=12)	0.02 mg/kg/day IV	393±123 (n=60 hr)	26.3±9.2	1.07 ± 0.20	0.038 ± 0.014
Mild Hepatic Impairment (n=6)	0.02 mg/kg/day IV (n=2) 7.7 mg PO	367 ± 107 (n=72 hr) 488 ± 320 (n=72 hr)	60.6 ± 43.8 Range: 27.8 - 141	3.1 ± 1.6	0.042 ± 0.02
Severe Hepatic Impairment (n=6, 5†)	0.02 mg/kg/day IV (n=4) 0.03 mg/kg/day IV (n=1) 5 mg PO (n=1) 4 mg PO (n=1)	762 ± 204 (n=120 hr) 299 ± 117 (n=144 hr) 608 (n=120 hr) 533 ± 156 (n=144 hr)	198 ± 158 Range:81 - 436 119 ± 35 Range: 85 - 178	3.9 ± 1.0 3.1 ± 3.4*	0.017 ± 0.013 0.016 ± 0.011*

* corrected for bioavailability
† 1 patient did not receive the PO dose

Patients with Renal Impairment

Tacrolimus pharmacokinetics have been determined in six patients with mild hepatic dysfunction (mean Pugh Score: 6.2) following IV administration of 0.02 mg/kg/day IV and oral administration of 7.7 mg/day PO. The mean clearance in patients with mild hepatic dysfunction was not substantially different from that in normal volunteers (see previous table). Tacrolimus pharmacokinetics were studied in 6 patients with severe hepatic dysfunction (mean Pugh score: > 10). The mean clearance was substantially lower in patients with severe hepatic dysfunction, irrespective of the route of administration *[see Dosage and Administration (2.5) and Use in Specific Populations (8.7)]*.

Racial or Ethnic Groups

The pharmacokinetics of tacrolimus have been studied following single IV and oral administration of tacrolimus to 10 African-American (12 ± 5 hr) and 10 Caucasian (11 ± 3 hr) healthy volunteers. There were no significant pharmacokinetic differences among the three ethnic groups following a 4-hour IV infusion of 0.015 mg/kg. However, after single oral administration of 5 mg, mean (±SD) tacrolimus C_{max} in African-Americans (23.6±12.1 ng/mL) was significantly lower than in Caucasians (49.2±13.8 ng/mL) and the Latino-Americans (38.2±15.8 ng/mL, p<0.01). Mean $AUC_{0-\infty}$ was significantly lower in African-Americans (203±115 ng•hr/mL) than Caucasians (344±188 ng•hr/mL) and Latino-Americans (274±180 ng•hr/mL). The mean (±SD) absolute oral bioavailability (F) in African-Americans (12±5%) and Latino-Americans (14±7.4%) was significantly lower than in Caucasians (19±9.8%, p<0.01). There was no significant difference in mean terminal $T_{1/2}$ among the three ethnic groups (range from 11 hours to 15 hours). The mean (±SD) oral clearance was significantly lower in African-Americans (0.12±0.06 L/hr/kg) than in Caucasians (0.19±0.09 L/hr/kg) and Latino-Americans (0.17±0.08 L/hr/kg). The mean (±SD) oral clearance in patients reported higher tacrolimus doses to attain similar trough concentrations *[see Dosage and Administration (2.2)]*.

Male and Female Patients

A formal trial to evaluate the effect of gender on tacrolimus pharmacokinetics has not been conducted, however, there was no difference in dosing by gender in the kidney transplant trial. A retrospective analysis of pharmacokinetics in healthy volunteers, and in kidney, liver and heart transplant patients indicated no gender-based differences.

Drug Interaction Studies

Frequent monitoring of whole blood concentrations and appropriate dosage adjustments of tacrolimus are recommended when concomitant use of the following drugs with tacrolimus is initiated or discontinued *[see Drug Interactions (7)]*.

- *Telaprevir*: In a single-dose study in 9 healthy volunteers, co-administration of tacrolimus (0.5 mg single dose) with telaprevir (750 mg three times daily for 11 days) increased the tacrolimus dose-normalized C_{max} by 9.3-fold and AUC by 70-fold compared to tacrolimus alone *[see Drug Interactions (7.2)]*.
- *Boceprevir*: In a single-dose study in 12 subjects, co-administration of tacrolimus (0.5 mg single dose) with boceprevir (800 mg three times daily for 11 days) increased tacrolimus C_{max} by 10 fold and AUC by 17-fold compared to tacrolimus alone *[see Drug Interactions (7.2)]*.
- *Nelfinavir*: Based on a clinical study of 5 liver transplant recipients, co-administration of tacrolimus with nelfinavir increased blood concentrations of tacrolimus significantly and, as a result, a reduction in the tacrolimus dose by an average of 16-fold was needed to maintain mean trough tacrolimus blood concentrations of 9.7 ng/mL. It is recommended to avoid concomitant use of tacrolimus and nelfinavir unless the benefits outweigh the risks *[see Drug Interactions (7.2)]*.
- *Rifampin*: In a study of 6 normal volunteers, a significant decrease in tacrolimus oral bioavailability (14±9% vs. 7±3%) was observed with concomitant rifampin administration (600 mg). In addition, there was a significant increase in tacrolimus clearance (0.053±0.017 L/hr/kg vs. 0.17±0.05 L/hr/kg). Tacrolimus dose requirements may be substantially increased *[see Drug Interactions (7.2)]*.
- *Magnesium/aluminum-hydroxide*: In a single-dose crossover study in healthy volunteers, co-administration of tacrolimus and magnesium-aluminum-hydroxide resulted in a 21% increase in the mean tacrolimus AUC and a 10% decrease in the mean tacrolimus C_{max} relative to tacrolimus administration alone *[see Drug Interactions (7.2)]*.
- *Ketoconazole*: In a study of 6 normal volunteers, a significant increase in tacrolimus oral bioavailability (14±5 vs. 30±8%) was observed with concomitant ketoconazole administration (200 mg). The apparent oral clearance of tacrolimus during ketoconazole administration was significantly decreased compared to tacrolimus alone (0.430±0.129 L/hr/kg vs. 0.066±0.018 L/hr/kg). The terminal $T_{1/2}$ of tacrolimus was prolonged (3.5±1.3 hr vs. 41.6±63.5 hr) and changed by ketoconazole co-administration, although it was highly variable between patients *[see Drug Interactions (7.2)]*.
- *Voriconazole* (see complete prescribing information for VFEND): Repeat oral dose administration of voriconazole (400 mg every 12 hours for one day, then 200 mg every 12 hours for 6 days) increased tacrolimus (0.1 mg/kg single dose) C_{max} and AUC_T in healthy subjects by an average of 2-fold (90% CI: 1.9, 2.5) and 3-fold (90% CI: 2.7, 3.8), respectively *[see Drug Interactions (7.2)]*.
- *Posaconazole* (see complete prescribing information for Noxafil): Repeat oral administration of posaconazole (400 mg twice daily for 7 days) increased tacrolimus (0.05 mg/kg single dose) C_{max} and AUC in healthy subjects by an average of 2-fold (90% CI: 2.07, 2.42) and 4.5-fold (90% CI 4.03, 5.19), respectively *[see Drug Interactions (7.2)]*.
- *Caspofungin* (see complete prescribing information for CANCIDAS): Caspofungin reduced the blood $AUC_{0-\tau}$ of tacrolimus by approximately 20%, peak blood concentration (C_{max}) by 16%, and 12-hour blood concentration (C_{12}) by 26% in healthy adult subjects when tacrolimus (2 doses of 0.1 mg/kg 12 hours apart) was administered on the 10th day of CANCIDAS 70 mg daily, as compared to results from a control period in which tacrolimus was administered alone *[see Drug Interactions (7.2)]*.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Carcinogenicity studies were conducted in male and female rats and mice. In the 80-week mouse oral study and in the 104-week rat oral study, no relationship of tumor incidence to tacrolimus dosage was found. The highest dose used in the mouse was 3.6 mg/kg/day (0.9 to 2.2 times the AUC at clinical doses of 0.075 to 0.2 mg/kg/day) and in the rat was 5 mg/kg/day (0.265 to 0.65 times the AUC at clinical doses of 0.075 to 0.2 mg/kg/day) *[see Boxed Warning and Warnings and Precautions (5.1)]*.

A 104-week dermal carcinogenicity study was performed in mice with tacrolimus ointment (0.03% - 3%), equivalent to tacrolimus doses of 1.1-116 mg/kg/day or 3.3-354 mg/m²/day. In the study, the incidence of skin tumors was minimal and the topical administration of tacrolimus did not result in systemic exposure sufficient to cause immunosuppression. There was, however, a significant elevation in the incidence of pleomorphic lymphoma in high-dose male (25/50) and female animals (27/50, and in the incidence of undifferentiated lymphoma in high-dose female animals (13/50) was noted in the mouse dermal carcinogenicity study. Lymphomas were noted in the mouse dermal carcinogenicity study at a daily dose of 3.5 mg/kg (0.1% tacrolimus ointment). No drug-related tumors were noted in the mouse dermal carcinogenicity study at a daily dose of 1.1 mg/kg (0.03% tacrolimus ointment). The relevance of topical administration of tacrolimus in the setting of systemic tacrolimus use is unknown.

The implications of these carcinogenicity studies to the human condition are limited; doses of tacrolimus were administered that likely induced immunosuppression in these animals, impairing their immune system's ability to inhibit emergent carcinogenesis.

Mutagenesis

No evidence of genotoxicity was seen in bacterial (*Salmonella* and *E. coli*) or mammalian (Chinese hamster lung-derived cells) *in vitro* assays of mutagenicity, the *in vitro* CHO/HGPRT assay of mutagenicity, or *in vivo* clastogenicity assays performed in mice; tacrolimus did not cause unscheduled DNA synthesis in rodent hepatocytes.

Impairment of Fertility

Tacrolimus, subcutaneously administered to male rats at paternally toxic doses of 2 mg/kg/day (1.6 to 4.3 times the recommended clinical dose range [0.2 to 0.075 mg/kg/day] on a mg/m² basis) or 3 mg/kg/day (2.4 to 6.4 times the recommended clinical dose range) resulted in a dose-related decrease in sperm count. Tacrolimus, administered orally at 1.6 mg/kg (0.8 to 2.2 times the clinical dose range) to male and female rats, prior to and during mating, as well as to dams during gestation and lactation, was associated with embryo/fetal and embryo/fetal and perinatal toxicity. Effects on female reproductive function (parturition) and embryo/fetal effects were indicated by a higher rate of pre- and post-implantation loss and increased numbers of undelivered females and stillborn pups at 1.6 mg/kg/day (0.8 to 2.2 times the clinical dose range on a mg/m² basis). Effects on male reproductive function (sperm count) and tacrolimus was associated with maternal and paternal toxicity as well as reproductive toxicity including marked adverse effects on estrus cycles, parturition, pup viability, and pup malformations.

14 CLINICAL STUDIES

14.1 Kidney Transplantation

Tacrolimus/Azathioprine (AZA)

The safety and efficacy of tacrolimus suppression in conjunction with azathioprine and corticosteroids following kidney transplantation was established in a randomized, multicenter, non-blinded, prospective trial. There were 412 kidney transplant patients enrolled at 19 clinical sites in the United States. Study therapy was initiated when renal function was stable as indicated by a serum creatinine ≤ 4 mg/dL (median of 4 days after transplantation, range 1 to 14 days). Patients were followed for 6 years of age were excluded.

There were 205 patients randomized to tacrolimus-based immunosuppression and 207 patients were randomized to cyclosporine-based immunosuppression. All patients received corticosteroids, induction therapy consisting of an antilymphocyte antibody preparation, corticosteroids, and azathioprine. Overall 1-year patient and graft survival was 96.1% and 89.6%, respectively.

Data from this trial of tacrolimus in conjunction with azathioprine indicate that during the first 3 months of that trial, 80% of the patients maintained trough concentrations between 7-20 ng/mL, and then between 5-15 ng/mL, through 1 year.

Tacrolimus/Mycophenolate mofetil (MMF)

Tacrolimus-based immunosuppression in conjunction with MMF, corticosteroids, and induction has been studied. In a randomized, open-label, multicenter trial (Study 1), 1589 kidney transplant patients received tacrolimus (Group C, n=401), sirolimus (Group D, n=399), or one of two cyclosporine (CsA) regimens (Group A, n=390 and Group B, n=399) in combination with MMF and corticosteroids. Induction therapy consisting of an interleukin-2 receptor antagonist was administered to 1578 patients. The trial was conducted outside the United States; the trial population was 83% Caucasian. In this study, mortality at 12 months in patients receiving tacrolimus/MMF was similar (3%) compared to patients receiving cyclosporine/M